

OXFORD

INTERNATIONAL
AQA EXAMINATIONS

INTERNATIONAL GCSE **COMBINED SCIENCE** **9204/CE**

Paper 2 Chemistry Extension

Mark scheme

June 2022

Version: 1.0 Final Mark Scheme



2 2 6 Y 9 2 0 4 / C E / M S

Mark schemes are prepared by the Lead Assessment Writer and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation events which all associates participate in and is the scheme which was used by them in this examination. The standardisation process ensures that the mark scheme covers the students' responses to questions and that every associate understands and applies it in the same correct way. As preparation for standardisation each associate analyses a number of students' scripts. Alternative answers not already covered by the mark scheme are discussed and legislated for. If, after the standardisation process, associates encounter unusual answers which have not been raised they are required to refer these to the Lead Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of students' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this mark scheme are available from [oxfordaqaexams.org.uk](https://www.oxfordaqaexams.org.uk)

Copyright information

OxfordAQA retains the copyright on all its publications. However, registered schools/colleges for OxfordAQA are permitted to copy material from this booklet for their own internal use, with the following important exception: OxfordAQA cannot give permission to schools/colleges to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Copyright © 2022 Oxford International AQA Examinations and its licensors. All rights reserved.

Level of response marking instructions

Level of response mark schemes are broken down into levels, each of which has a descriptor. The descriptor for the level shows the average performance for the level. There are marks in each level.

Before you apply the mark scheme to a student's answer read through the answer and annotate it (as instructed) to show the qualities that are being looked for. You can then apply the mark scheme.

Step 1 Determine a level

Start at the lowest level of the mark scheme and use it as a ladder to see whether the answer meets the descriptor for that level. The descriptor for the level indicates the different qualities that might be seen in the student's answer for that level. If it meets the lowest level then go to the next one and decide if it meets this level, and so on, until you have a match between the level descriptor and the answer. With practice and familiarity you will find that for better answers you will be able to quickly skip through the lower levels of the mark scheme.

When assigning a level you should look at the overall quality of the answer and not look to pick holes in small and specific parts of the answer where the student has not performed quite as well as the rest. If the answer covers different aspects of different levels of the mark scheme you should use a best fit approach for defining the level and then use the variability of the response to help decide the mark within the level, ie if the response is predominantly level 3 with a small amount of level 4 material it would be placed in level 3 but be awarded a mark near the top of the level because of the level 4 content.

Step 2 Determine a mark

Once you have assigned a level you need to decide on the mark. The descriptors on how to allocate marks can help with this. The exemplar materials used during standardisation will help. There will be an answer in the standardising materials which will correspond with each level of the mark scheme. This answer will have been awarded a mark by the Lead Examiner. You can compare the student's answer with the example to determine if it is the same standard, better or worse than the example. You can then use this to allocate a mark for the answer based on the Lead Examiner's mark on the example.

You may well need to read back through the answer as you apply the mark scheme to clarify points and assure yourself that the level and the mark are appropriate.

Indicative content in the mark scheme is provided as a guide for examiners. It is not intended to be exhaustive and you must credit other valid points. Students do not have to cover all of the points mentioned in the Indicative content to reach the highest level of the mark scheme.

An answer which contains nothing of relevance to the question must be awarded no marks.

Information to Examiners

1. General

The mark scheme for each question shows:

- the marks available for each part of the question
- the total marks available for the question
- the typical answer or answers which are expected
- extra information to help the Examiner make his or her judgement
- the Assessment Objectives, level of demand and specification content that each question is intended to cover.

The extra information is aligned to the appropriate answer in the left-hand part of the mark scheme and should only be applied to that item in the mark scheme.

At the beginning of a part of a question a reminder may be given, for example: where consequential marking needs to be considered in a calculation; or the answer may be on the diagram or at a different place on the script.

In general the right-hand side of the mark scheme is there to provide those extra details which confuse the main part of the mark scheme yet may be helpful in ensuring that marking is straightforward and consistent.

2. Emboldening and underlining

- 2.1** In a list of acceptable answers where more than one mark is available ‘any **two** from’ is used, with the number of marks emboldened. Each of the following bullet points is a potential mark.
- 2.2** A bold **and** is used to indicate that both parts of the answer are required to award the mark.
- 2.3** Alternative answers acceptable for a mark are indicated by the use of **or**. Different terms in the mark scheme are shown by a / ; eg allow smooth/free movement.
- 2.4** Any wording that is underlined is essential for the marking point to be awarded.

3. Marking points

3.1 Marking of lists

This applies to questions requiring a set number of responses, but for which students have provided extra responses. The general principle to be followed in such a situation is that ‘right + wrong = wrong’.

Each error/contradiction negates each correct response. So, if the number of errors/contradictions equals or exceeds the number of marks available for the question, no marks can be awarded.

However, responses considered to be neutral (indicated as * in example 1) are not penalised.

Example 1: What is the pH of an acidic solution?

[1 mark]

Student	Response	Marks awarded
1	green, 5	0
2	red*, 5	1
3	red*, 8	0

Example 2: Name two planets in the solar system.

[2 marks]

Student	Response	Marks awarded
1	Neptune, Mars, Moon	1
2	Neptune, Sun, Mars, Moon	0

3.2 Use of chemical symbols/formulae

If a student writes a chemical symbol/formula instead of a required chemical name, full credit can be given if the symbol/formula is correct and if, in the context of the question, such action is appropriate.

3.3 Marking procedure for calculations

Marks should be awarded for each stage of the calculation completed correctly, as students are instructed to show their working. Full marks can, however, be given for a correct numerical answer, without any working shown.

3.4 Interpretation of 'it'

Answers using the word 'it' should be given credit only if it is clear that the 'it' refers to the correct subject.

3.5 Errors carried forward

Any error in the answers to a structured question should be penalised once only.

Papers should be constructed in such a way that the number of times errors can be carried forward is kept to a minimum. Allowances for errors carried forward are most likely to be restricted to calculation questions and should be shown by the abbreviation ecf in the marking scheme.

3.6 Phonetic spelling

The phonetic spelling of correct scientific terminology should be credited **unless** there is a possible confusion with another technical term.

3.7 Brackets

(.....) are used to indicate information which is not essential for the mark to be awarded but is included to help the examiner identify the sense of the answer required.

3.8 Allow

In the mark scheme additional information, 'allow' is used to indicate creditworthy alternative answers.

3.9 Ignore

Ignore is used when the information given is irrelevant to the question or not enough to gain the marking point. Any further correct amplification could gain the marking point.

3.10 Do not accept

Do **not** accept means that this is a wrong answer which, even if the correct answer is given as well, will still mean that the mark is not awarded.

Question	Answers	Extra information	Mark	AO/Spec. Ref.
01.1	rubidium		1	AO1 3.13.1a
01.2	chlorine		1	AO2 3.7.2i
01.3	helium		1	AO1 3.7.3c
01.4	potassium	allow K	1	AO2 3.8.1c
01.5	36		1	AO2 3.8.1e
01.6	(their atoms) have stable arrangements of electrons	allow (their atoms) have full outer energy levels allow (their atoms) have full outer shells	1	AO1 3.7.3c
01.7	(conversion 1.50 kg =) 1500 g	allow correct use of incorrectly / not converted values from step 1	1	AO2 × 3 3.8.1d
	(solubility =) $43.2 \times \frac{100}{1500}$		1	
	= 2.88 (mol/100 g)		1	
Total			9	

Question	Answers	Extra information	Mark	AO/Spec. Ref.
02.1	$4\text{NH}_3 + 5 \text{O}_2 \rightarrow 4\text{NO} + 6\text{H}_2\text{O}$	allow multiples	1	AO2 3.12.1a
02.2	gain of oxygen by ammonia / nitrogen / hydrogen	allow 1 mark for gain of oxygen	2	AO1 AO2 3.9.1.1d
02.3	activation energy		1	AO1 3.15.2a
02.4	particles have more energy (so) when the particles collide there is enough energy to react or (so) when the particles collide the energy is greater than / equal to the activation energy alternative approach: more particles have energy greater than / equal to the activation energy (1) (so) particles react when they collide (1)		1 1	AO1 × 2 3.14.1b
02.5	(level of) products is below (level of) reactants	allow the energy decreases (overall) allow energy is transferred to the surroundings ignore references to bond making / breaking	1	AO3 3.15.2a
02.6	(a catalyst) increases the rate of the reaction (by) providing a different reaction pathway or (by) lowering the activation energy	allow (a catalyst) speeds up the reaction	1 1	AO1 × 2 3.14.1h 3.15.2e

02.7	low boiling point non-conductor of electricity		1 1	AO1 × 2 3.8.2ce
02.8	$(M_r = 14 + (16 \times 2) =) 46$ $(\% \text{ by mass} =) \left(\frac{14}{46}\right) \times 100$ $= 30.43$ $= 30.4 (\%)$	allow correct use of an incorrectly calculated value of M_r allow 30.43478261 allow a correctly calculated answer given to 3 significant figures from an incorrect attempt at the percentage by mass of nitrogen	1 1 1 1	AO2 × 4 3.12.2b
Total			15	

Question	Answers	Extra information	Mark	AO/Spec. Ref.
03.1	CaCO ₃		1	AO2 × 2
	(s)		1	3.11.1e 3.12.1a 3.12.1b

Question	Answers	Mark	AO/Spec. Ref.
----------	---------	------	---------------

03.2	Level 3: The design/plan would lead to the production of a valid outcome. All key steps are identified and logically sequenced.	5–6	AO4 × 6 3.9.1.2a
	Level 2: The design/plan would not necessarily lead to a valid outcome. Most steps are identified, but the plan is not fully logically sequenced.	3–4	
	Level 1: The design/plan would not lead to a valid outcome. Some relevant steps are identified, but links are not made clear.	1–2	
	No relevant content.	0	
	Indicative content • add known mass of (Group 2) carbonate to boiling tube • heat the (Group 2) carbonate • measure time for limewater to change using stop clock or use of alternative methods such as counting bubbles or comparison of loss of mass of boiling tube and contents • repeat using other (Group 2) carbonates • same mass of each (Group 2) carbonate • same sized pieces of each (Group 2) carbonate • fresh limewater for each experiment • same volume of limewater • judge same degree of cloudiness of limewater such as time for 'x' to disappear • repeat each experiment • the greater the time the slower the rate of reaction		

Total			8
--------------	--	--	----------

Question	Answers	Extra information	Mark	AO/Spec. Ref.
04.1	fractional distillation		1	AO1 3.16.1.1c
04.2	boiling point	allow temperature at which the hydrocarbons evaporate / condense	1	AO1 3.16.1.1c
04.3	C ₈ H ₁₆	allow 1 mark for $\frac{112}{14}$ or 8 carbon atoms per molecule	2	AO2 × 2 3.12.2c
04.4	any two from: • high boiling point or low volatility • low flammability • burn with a smoky flame.		2	AO1 × 2 3.16.1.2c
04.5	CO C	ignore names	1 1	AO1 AO2 3.16.1.2d
04.6	coal contains sulfur (so) sulfur dioxide is produced (when coal burns) (and) sulfur dioxide dissolves in water to form acid rain	allow (so) oxides of nitrogen are produced allow (and) oxides of nitrogen dissolve in water to form acid rain	1 1 1	AO1 AO2 AO3 3.16.1.2d
Total			11	

Question	Answers	Extra information	Mark	AO/Spec. Ref.
05.1	(chlorine) to produce bleach or to produce plastics	allow to sterilise water	1	AO1 × 2 3.9.2j
	(sodium hydroxide) to produce soap		1	
05.2	colour change from green to blue / purple		1	AO3 × 3 3.9.2j 3.11.1f
	(because) sodium chloride solution is neutral	allow (because) sodium chloride solution has a pH of 7	1	
	(and because) sodium hydroxide solution is alkaline	allow (and because) sodium hydroxide solution has a pH above 7	1	
05.3	sodium is more reactive than hydrogen		1	AO2 3.9.2g
05.4	$2\text{H}^+ + 2\text{e}^- \rightarrow \text{H}_2$	allow 1 mark for $\text{H}^+ + \text{e}^- \rightarrow \text{H}_2$ regardless of any balancing numbers	2	AO2 × 2 3.9.2f
05.5	giant structure (of carbon atoms)		1	AO1 x4 3.8.3c
	covalent bonding		1	
	each (carbon) atom forms 3 (covalent) bonds		1	
	forming layers (of atoms)		1	


05.6	$(M_r = 35.5 \times 2) = 71$ $(\text{moles} =) \frac{0.0213}{71}$ $= 0.00030$ $= 3.0 \times 10^{-4} \text{ (mol)}$	allow correct use of an incorrect value of M_r allow a correctly calculated and rounded conversion to standard form of an incorrect calculation of number of moles	1 1 1 1	AO2 × 4 3.12.3a
Total			16	

Question	Answers	Extra information	Mark	AO/Spec. Ref.
06.1	(moles = $0.150 \times 0.250 =$) 0.0375 ($M_r = 39 + 16 + 1 =$) 56 (mass =) 0.0375×56 = 2.1 (g) alternative approach: ($M_r = 39 + 16 + 1 =$) 56 (1) (mass per $\text{dm}^3 = 0.150 \times 56 =$) 8.4 ($\text{g}/\text{dm}^3$) (1) (mass =) 8.4×0.250 (1) = 2.1 (g) (1)	allow correct use of an incorrectly calculated value of M_r and / or incorrectly calculated value of moles allow correct use of an incorrectly calculated value of M_r	1 1 1 1	AO2 $\times$ 4 3.12.3a 3.12.4a
06.2	ionises completely (in aqueous solution)		1	AO3 3.16.3.2b
06.3	potassium sulfate		1	AO2 3.11.1c

06.4	half the volume of sulfuric acid needed to neutralise the alkali compared with hydrochloric acid	allow 14.5 (cm ³) sulfuric acid and 29.0 (cm ³) hydrochloric acid	1	AO3 × 3 3.11.1f 3.12.4c
	(because) sulfuric acid produces twice as many H ⁺ ions	allow sulfuric acid neutralises twice the number of moles of alkali	1	
	per mole / molecule of acid	if no other mark awarded, allow 1 mark for a smaller volume of sulfuric acid because it produces more H ⁺ ions	1	
Total			9	

Question	Answers	Extra information	Mark	AO/Spec. Ref.
07.1	$ \begin{array}{ccccccc} & \text{H} & & \text{H} & & \text{H} & \\ & & & & & & \\ \text{C} & = & \text{C} & - & \text{C} & - & \text{C} & - & \text{H} \\ & & & & & & \\ & \text{H} & & \text{H} & & \text{H} & & \text{H} \end{array} $ or $ \begin{array}{ccccccc} & \text{H} & & & & \text{H} & & \text{H} \\ & & & & & & & \\ \text{H} & - & \text{C} & - & \text{C} & = & \text{C} & - & \text{C} & - & \text{H} \\ & & & & & & & \\ & & \text{H} & & & & & \text{H} \end{array} $	allow 1 mark for C ₄ H ₈	2	AO3 × 2 3.8.3a 3.16.1.3b 3.16.1.3c
07.2	vaporises	allow turns into a gas allow evaporates	1	AO4 3.16.1.3a
07.3	C ₈ H ₁₈ is a liquid (when cooled)	allow C ₈ H ₁₈ is not a gas allow C ₈ H ₁₈ condenses (when cooled)	1	AO3 3.16.1.3a
07.4	C ₂ H ₆	allow 1 mark for a formula with 6 hydrogen atoms	2	AO3 × 2 3.16.1.2e 3.16.1.3b 3.16.1.3d
Total			6	

Question	Answers	Extra information	Mark	AO/Spec. Ref.
08.1	boiling point increases down the group		1	AO1 AO2 × 2 3.8.2d 3.13.1d
	(because) size of molecules increases down the group		1	
	(so) more energy is required to overcome the intermolecular forces	allow (so) the strength of the intermolecular forces increases	1	
08.2	$\text{Br}_2 + 2\text{KI} \rightarrow 2\text{KBr} + \text{I}_2$	allow 1 mark for $\text{Br}_2 + \text{KI} \rightarrow \text{KBr} + \text{I}_2$ regardless of balancing numbers	2	AO2 × 2 3.12.1a 3.13.1e
08.3	solution turns from orange to colourless	allow solution turns from orange to brown allow colour of solution will fade	1	AO3 × 2 3.8.1e
	(purple-black) solid forms	ignore iodine forms	1	
08.4	the outer electrons (in bromine) are closer to the nucleus	allow converse allow (bromine has) a smaller atom allow (bromine has) fewer shells allow (bromine has) fewer energy levels	1	AO1 × 3 3.13.1f
	(so outer electrons are) more strongly attracted to the nucleus	allow (so) less shielding	1	
	(and so) an additional electron is more easily gained		1	

08.5	1 pair of bonded electrons 6 non-bonded electrons on Br and 6 non-bonded electrons on F	allow any combination of circles, dots, crosses or e ⁽⁻⁾ an answer of  scores 2 marks	1 1	AO2 × 2 3.8.1g
Total			12	

Question	Answers	Extra information	Mark	AO/Spec. Ref.
09.1	14		1	AO2 3.7.2cf
09.2	(magnesium contains) delocalised electrons	allow free electrons for delocalised electrons	1	AO1 × 3 3.8.1j 3.8.2g
	from the highest occupied energy level	allow outer shell for energy level	1	
	(so) electrons can move through the structure / metal	allow electrons carry (thermal) energy through the structure	1	
09.3	Mg ₂ SiO ₄		1	AO2 3.8.1c
09.4	magnesium is more reactive than carbon		1	AO1 × 2 AO2 3.9.1.1a 3.9.1.1e
	(so) is extracted by electrolysis (of a molten compound)		1	
	(which) requires large amounts of energy / electricity		1	
09.5	($M_r = 24 + 12 + 48 =$) 84		1	AO2 × 4 3.12.2a 3.12.3a
	$\left(\text{moles} = \frac{2100}{84} =\right)$ 25	allow correct use of an incorrectly calculated value of M_r	1	
	(mass =) 25 × 24	allow correct use of an incorrectly calculated value of moles	1	
	= 600 (g)		1	
	alternative approach: ($M_r = 24 + 12 + 48 =$) 84 (1) (proportion of magnesium in magnesium carbonate =) $\frac{24}{84}$ (1) 0.2857 × 2100 (1) = 600 (g) (1)	allow correct use of an incorrectly calculated value of M_r		

09.6	D = barium sulfate	allow BaSO_4 do not accept magnesium sulfate do not accept calcium sulfate	1	AO3 × 3 3.10.2b 3.10.2f
	E = aluminium hydroxide	allow $\text{Al}(\text{OH})_3$	1	
Total			14	