



Cambridge International AS & A Level

CANDIDATE
NAMECENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

MATHEMATICS

9709/25

Paper 2 Pure Mathematics 2

May/June 2026

1 hour 15 minutes

You must answer on the question paper.

You will need: List of formulae (MF19)

INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen. Do **not** use correction fluid or tape.
- Do **not** write on any bar codes.
- If additional space is needed, you should use the lined page at the end of this booklet; the question number or numbers must be clearly shown.
- You should use a calculator where appropriate.
- You must show all necessary working clearly; no marks will be given for unsupported answers from a calculator.
- Give non-exact numerical answers correct to 3 significant figures, or 1 decimal place for angles in degrees, unless a different level of accuracy is specified in the question.

INFORMATION

- The total mark for this paper is 50.
- The number of marks for each question or part question is shown in brackets [].

This document has **16** pages. Any blank pages are indicated.



1 Find $\int_a^{2a} (1 + e^{-x})^2 dx$. Give your answer in terms of the positive constant a .

[4]

[illegible]




[4]

[illegible]



3



The variables x and y satisfy the equation $y = ka^{x+2}$, where k and a are positive constants. The graph of $\ln y$ against x is a straight line passing through the points $(1.5, 3.22)$ and $(4.0, 4.69)$, as shown in the diagram.

Find the values of k and a correct to 2 significant figures.

[5]

PapaCambridge
papacambridge.com





- 4 A curve has equation $y = \frac{e^{2x}}{e^x + 2}$. The curve crosses the y -axis at the point P .

Find an equation of the normal to the curve at P . Give your answer in the form $ax + by + c = 0$, where a , b and c are integers. [6]

This image shows a full page of blank handwriting practice paper. It features multiple rows of horizontal lines. Each row consists of a solid top line, a dashed midline, and a solid bottom line, providing a guide for letter height and placement. The paper is white with light gray or blue lines. There are faint, diagonal watermarks across the page that read "PapaCambridge.com".


- 5 (a) Solve the inequality $|4x - 7| \leq |2x + 3|$.

[4]

- (b) Hence, find the smallest and largest integers, N , that satisfy the inequality

$$\left| 4 \ln\left(\frac{1}{2}N\right) - 7 \right| \leq \left| 2 \ln\left(\frac{1}{2}N\right) + 3 \right|.$$

[3]



The diagram shows the curve with equation $y = f(x)$, where

$$f(x) = ax^4 + 4x^3 - 3ax^2 - 21x - 18$$

and a is a constant. It is given that $(x + 2)$ is a factor of $f(x)$.

(a) Find the value of a .

[2]

.....

.....

.....

.....

.....

.....

.....

(b) Show that the positive root of the equation $f(x) = 0$ satisfies the equation

$$x = \sqrt[3]{3x + 4.5}.$$

[3]

.....

.....

.....

.....

.....

.....

.....






- (c) Use an iterative formula based on the equation $x = \sqrt[3]{3x+4.5}$ to find the positive root of the equation $f(x) = 0$ correct to 4 significant figures. Give the result of each iteration to 6 significant figures.

[3]





The diagram shows part of the curve with equation $y = 8 \cos 2x - \frac{1}{\cos^2 2x}$. The shaded region is bounded by the curve and the x -axis.

Find the exact area of the shaded region.


[7]

[illegible]



This image shows a full page of blank handwriting practice paper. It features multiple sets of horizontal lines. Each set consists of a solid top line, a dashed midline, and a solid bottom line, providing a guide for letter height and placement. The background is white, and the lines are light gray. Faint, diagonal watermarks reading "PapaCambridge.com" are repeated across the entire page.





The diagram shows the curve with parametric equations

$$x = 6t - \sin 2t, \quad y = 3 \cos 2t + 5 \sin t,$$

for $0 \leq t < \frac{1}{2}\pi$. The curve has a maximum point, M .

- (a)** Find the gradient of the curve at the point for which $t = 0$.

[4]



(b) Find the exact y -coordinate of M .

[5]



[illegible]



DO NOT WRITE IN THIS MARGIN





Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (Cambridge University Press & Assessment) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced online in our Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download at www.cambridgeinternational.org after the live examination series.

Cambridge International Education is the name of our awarding body and a part of Cambridge University Press & Assessment, which is a department of the University of Cambridge.

